

Physical Therapy Prescription – ACL Reconstruction with Hamstring Autograft

Procedure: ACLR with hamstring tendon autograft

Frequency: 2-3 times per week for 6 weeks

PHASE I (Weeks 0 – 4): Period of protection, decrease edema, activate quadriceps

- **Weightbearing:** As tolerated with crutches, with goal of discontinuing crutches by 10 days (may be modified if meniscus repair/transplant or articular cartilage surgery performed at time of ACLR)
- **Hinged Knee Brace:**
 - Locked in full extension for ambulation and sleeping (weeks 0-4)
 - i. Only unlock when cleared by practice
- **Range of Motion:** AAROM → AROM as tolerated
- **Therapeutic Exercises:** Patellar mobs, quad/hamstring sets, heel slides, non-weightbearing Gastroc/Soleus stretching, straight-leg raises with brace in full extension until quad strength prevents extension lag
- **Modalities:** Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)
- **Precautions:** No hamstring stretching until 4 weeks postop

Phase II (Weeks 4 – 6)

- **Weightbearing:** As tolerated, unassisted
- **Hinged Knee Brace:** Discontinue once full extension achieved with no evidence of extension lag
- **Range of Motion:** Maintain full knee extension, work on progressive knee flexion
- **Therapeutic Exercises:** Patellar mobs, quad/hamstring sets, closed chain extension exercises, hamstring curls, toe raises, balance exercises, progress to weightbearing Gastroc/Soleus stretch
 - Begin use of the stationary bicycle
- **Modalities:** Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase III (Weeks 6 – 16)

- **Range of Motion:** Full, painless
- **Therapeutic Exercises:** Advance closed chain strengthening exercises and proprioception activities
 - Begin use of the Stairmaster/Elliptical at **8 weeks**
 - Straight ahead running permitted at **12 weeks**
- **Modalities:** Per therapist, including electrical stimulation, ultrasound, heat (before), ice (after)

Phase IV (Weeks 16 – 24): Gradual return to athletic activity

- **16 weeks:** begin jumping
- **20 weeks:** advance to sprinting, backward running, cutting/pivoting/changing direction
- **24 weeks:** advance as tolerated, consider **functional sports assessment**

Phase V (>6 months): Gradual return to athletic activity

- Gradual return to sports participation after completion of functional sports assessment
- Encourage maintenance program based off functional sports assessment

Signature: _____

Date: _____