

**Kade McQuivey, MD**  
Department of Orthopedics  
Sports Medicine and Shoulder Surgery  
Office Phone: 623.876.3870  
Fax: 623.337.9279  
[www.KadeMcQuiveyMD.com](http://www.KadeMcQuiveyMD.com)



Banner New River Trails  
Sun City West Medical Center  
Thunderbird Medical Center

## **Rehabilitation Protocol**

### **Acute Coracoclavicular Ligament Reconstruction**

#### **Phase I: Protect Repair** *(0 to 6 weeks after surgery)*

- Patients may shower immediately over clear plastic, waterproof dressing
- Sutures are all underneath the skin and will dissolve on their own
- Ice or cold flow systems encouraged for the first week at a minimum: should be used 3-4 times per day.
- Sling should be worn at all times.
- Initiate exercise program:  
    Immediate elbow, forearm and hand range of motion out of sling
- No lifting with involved extremity.
- AVOID scapular ROM exercises.

#### **Phase II: Progress ROM & Protect Repair** *(6 to 12 weeks after surgery)*

- May discontinue sling.
- Pendulum exercises
- Passive and active assistive ER at the side to 30, flexion to 130
- Initiate gentle rotator cuff strengthening.
- Initiate scapular AROM exercises.

#### **Phase III: Full Function** *(3 months after surgery)*

- Advance active and passive ROM in all planes to tolerance.
- Advance rotator cuff and scapular stabilizer strengthening.
- Initiate functional progression to sports specific activities at 5 months.
- Discontinue lifting restrictions after 5 months