

## Rehabilitation Protocol Triceps Tendon Repair

### Phase I: Protect Repair (0 to 2 weeks)

- **Precautions**
  - Elbow immobilized in posterior splint at 60° elbow flexion for 1st 2 weeks
  - No active elbow extension 6 weeks
- **ROM**
  - Gradual ↑ Active/Passive ROM of shoulder in all planes while in splint
  - Wrist/hand/finger full AROM in splint
- **Strength**
  - Scapular retractions
  - Shoulder shrugs
- **Modalities**
  - Hot pack before treatment
  - E-stim, TENS as needed
  - Ice 10-15 minutes after treatment
- **Goals of Phase I**
  - Control pain and inflammation
  - Protect repair
  - Independent in HEP

### Phase II: Progress Protected ROM and Function (3-6 weeks)

- **Precautions**
  - Elbow placed in a *hinged ROM brace* allowing 30°-60°
  - *Brace* to be worn at all times except during exercise or bathing
  - *Passive* ROM ONLY for elbow extension
- **ROM**

**Kade McQuivey, MD**  
Department of Orthopedics  
Sports Medicine and Shoulder Surgery  
Office Phone: 623.876.3870  
Fax: 623.337.9279  
[www.KadeMcQuiveyMD.com](http://www.KadeMcQuiveyMD.com)



Banner New River Trails  
Sun City West Medical Center  
Thunderbird Medical Center

- Hinged Brace Range of Motion Progression (ROM progression may be adjusted base on Surgeon's assessment of the surgical repair.)
- Week 2-3: 30°-60°
- Week 4-5: 15°-90°
- Week 6-7: 10°-110°
- Week 8: 0°-125°
- Forearm: Initiate AAROM pronation and supination
- Progress to active pronation and supination (wk 4)
- Shoulder AROM as needed in brace
  
- **Strength (in brace)**
  - Isometric shoulder exercises
  - Supine/standing rhythmic stabilizations
  - Wrist/hand: grip strengthening
  - Standing flexion and scaption
  - Side-lying ER
  - Isometric biceps pain free (week 6)
  
- **Manual**
  - Scar mobilization
  - Passive elbow extension
  - Joint mobs as needed
  
- **Modalities**
  - Heat/hot pack before therapy
  - US to incision as needed
  - Ice 10-15 minutes
  
- **Goals of Phase II**
  - Protection of repair
  - Gradual increase in ROM
  - Initiate strengthening to surrounding tissues
    - Improve scapular stability

**Kade McQuivey, MD**  
Department of Orthopedics  
Sports Medicine and Shoulder Surgery  
Office Phone: 623.876.3870  
Fax: 623.337.9279  
[www.KadeMcQuiveyMD.com](http://www.KadeMcQuiveyMD.com)



Banner New River Trails  
Sun City West Medical Center  
Thunderbird Medical Center

### **Phase III: Reach full ROM** (*7-12 weeks*)

- **ROM**
  - Week 8 progress to full ROM of elbow; discontinue brace if adequate motor control
  - Initiate UBE light resistance
  - Exercises
  - Ball roll outs on table
  - Wall walk
  - Pulley
  
- **Strength**
  - Tricep/elbow extension progression
  - 6 weeks: initiate AROM
  - 8 weeks: initiate light Theraband resistance
  - Theraband IR/ER shoulder
  - Theraband bicep extension
  - Prone dumbbell Therex
  - Rhythmic stabilization
  
- **Manual**
  - Passive elbow extension if lacking
  - Joint mobs as needed to regain full flexion
  - Week 10: Passive or contract relax to gain full flexion if still lacking

### **Phase IV: Full ROM and Strength** (*>12 weeks*)

- **Strength**
  - Progress strengthening program with increase in resistance and high speed repetition
  - Bicep curls with dumbbells
  - Initiate IR/ER exercises at 90° abduction
  - Progress rhythmic stabilization activities to include standing PNF patterns with tubing
  - Initiate plytoss – double arm progress to single arm

**Kade McQuivey, MD**  
Department of Orthopedics  
Sports Medicine and Shoulder Surgery  
Office Phone: 623.876.3870  
Fax: 623.337.9279  
[www.KadeMcQuiveyMD.com](http://www.KadeMcQuiveyMD.com)



Banner New River Trails  
Sun City West Medical Center  
Thunderbird Medical Center

- Initiate sport specific drills and functional activities
- Initiate interval throwing program week 16-20
- Initiate light upper body plyometric program week 16-20
- Progress isokinetics to 90° abduction at high speeds
  
- **Modalities**
  - Ice 15-20 minutes
  
- **Goals of Phase IV**
  - Full painless ROM
  - Maximize upper extremity strength and endurance
  - Maximize neuromuscular control
  - Optimize shoulder mechanics/kinematics
  - Optimize core stability
  - Initiate sports specific training/functional training